



An Evaluation of Pharmacy Data for Surveillance of Gastrointestinal and Respiratory Outbreaks

Yevgeniy Elbert¹, MS Shilpa Hakre¹, MPH, DrPH Howard Burkom², PhD Julie Pavlin¹, MD, MPH

November 15, 2004

¹Walter Reed Army Institute of Research, Silver Spring, MD ²Johns Hopkins Applied Physics Laboratory, Laurel, MD

maintaining the data needed, and of including suggestions for reducing	llection of information is estimated to completing and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding ar OMB control number.	ion of information. Send comments arters Services, Directorate for Information	regarding this burden estimate mation Operations and Reports	or any other aspect of th , 1215 Jefferson Davis l	is collection of information, Highway, Suite 1204, Arlington		
1. REPORT DATE 15 NOV 2004		2. REPORT TYPE N/A		3. DATES COVE	RED		
4. TITLE AND SUBTITLE					5a. CONTRACT NUMBER		
An Evaluation of Pharmacy Data for Surveillance of Gastrointestinal and					5b. GRANT NUMBER		
Respiratory Outbreaks				5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)					5d. PROJECT NUMBER		
					5e. TASK NUMBER		
				5f. WORK UNIT NUMBER			
	ZATION NAME(S) AND AD TRACE	D	8. PERFORMING ORGANIZATION REPORT NUMBER				
9. SPONSORING/MONITO	RING AGENCY NAME(S) A	10. SPONSOR/MONITOR'S ACRONYM(S)					
					11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAIL Approved for publ	LABILITY STATEMENT ic release, distributi	on unlimited					
	otes 49, 2004 Scientific C land on 15-17 Nove		0				
14. ABSTRACT							
15. SUBJECT TERMS							
16. SECURITY CLASSIFIC	17. LIMITATION OF	18. NUMBER	19a. NAME OF				
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	ABSTRACT UU	OF PAGES 40	RESPONSIBLE PERSON		

Report Documentation Page

Form Approved OMB No. 0704-0188

ESSENCE

Electronic Surveillance System for the Early Notification of Community-based Epidemics


- A U.S. Department of Defense (DoD) system
- Designed to detect infectious disease outbreaks
- Serves military active duty members, their beneficiaries, and retirees
- Uses mainly ICD-9-CM codes from outpatient visits
- Delay of 1- 4 days from patient visit date to data capture date by ESSENCE

Pharmacy Data Transaction Service (PDTS)

- Developed by the DoD Pharmacoeconomic Center (PEC) program, managed by WebMD
- Data repository for prescriptions filled by beneficiaries of the military health system
- Approx. 61.1 million prescriptions filled annually¹
 - 82% of prescriptions from MTFs (n=587)
 - 18% from:
 - Tricare Managed Care Support Contractors Network pharmacies (n=50,000)
 - National Mail Order Pharmacy (n=1)

ESSENCE





Update every 24 hours



Tricare pharmacy transactions

3.2 seconds

PDTS Fields

- The PDTS formulary and standards maintained through WebMD's subscription to the American Hospital Formulary Service (AHFS), the National Council for Prescription Drug Programs (NCPDP), and First Data Bank, Inc.
- Medication id fields in PDTS
 - Drug name (Label)
 - Therapeutic class number (GCN)
 - Therapeutic class code (GC3)
 - Other
- Previous study by Scott Eader et al. using ESSENCE data for the DC metropolitan area
 - Significant positive correlation between outpatient visit and PDTS data for GC3 codes identified for Respiratory (RESP) and gastrointestinal (GI) syndromes
- GC3: developed by First Data Bank, Inc.
 - 3 characters (alpha, numeric, alpha)
 - organ system, pharmacological class, specific therapeutic class, respectively

PDTS Commonly filled medications, 2003

GC3	DESCRIPTION	COUNT	PERCENT
S2B	NSAIDS, CYCLOOXYGENASE INHIBITOR - TYPE	5603369	Painkillers
НЗА	ANALGESICS,NARCOTICS	4256802	4.81 Fallikillers
M4E	LIPOTROPICS	4049267	4.58
Z2A	ANTIHISTAMINES	3812841	4.34 Cholestrol
D4K	GASTRIC ACID SECRETION REDUCERS	3395463	3.84 reducers
A4D	HYPOTENSIVES, ACE INHIBITORS	2659102	3.01
J7C	BETA-ADRENERGIC BLOCKING AGENTS	2373543	2.68
H2S	SELECTIVE SEROTONIN REUPTAKE INHIBITOR (SSRIS)	2230604	2.52 Allergy
W1A	PENICILLINS	2039720	2.31 relievers
A9A	CALCIUM CHANNEL BLOCKING AGENTS	1966936	2.22
P3A	THYROID HORMONES	1700412	1.92
B3J	EXPECTORANTS	1654859	1.87
J5D	BETA-ADRENERGIC AGENTS	1557748	1.76
Q7P	NASAL ANTI-INFLAMMATORY STEROIDS	1420677	1.61
H2F	ANTI-ANXIETY DRUGS	1311393	1.48
H4B	ANTICONVULSANTS	1299466	1.47
P5A	GLUCOCORTICOIDS	1265605	1.43
R1F	THIAZIDE AND RELATED DIURETICS	1247417	1.41
W1D	MACROLIDES	1242950	1.41
A4F	HYPOTENSIVES,ANGIOTENSIN RECEPTOR ANTAGONIST	1223731	1.38

STUDY OBJECTIVES

- Determine medications commonly prescribed for GI and RESP syndromes during outbreaks
- Examine trends in daily counts of medications filled for GI, RESP, Asthma visits during outbreaks
- 3) Conduct retrospective surveillance on drug groups that correlated with GI and RESP outpatient visits during outbreaks


METHODS

- List compiled of known GI and RESP outbreaks
- Commonly prescribed non-refill medications identified for each outbreak
 - By linking ambulatory and pharmacy data by date of visit/prescription written and unique encrypted identifiers common to both data sets
- Trends in medications filled and GI/Resp visits investigated during outbreaks
 - Moving 7-day averages of daily counts

RESULTSOutbreaks


<u>Syndrome</u>	No. of Outbreaks	<u>Viral</u>	<u>Bacterial</u>	<u>Unknown</u>
GI	21	13	2	6
RESP	7	2	4	1
Total	28	15	6	7

GI Outbreaks




- Food and water samples from the FTX tested
 - 5/13 samples tested positive for Campylobacter jejuni

Top 5 ICD-9 codes used during a C. jejuni outbreak




Top 5 prescriptions written during C. jejuni outbreak



RESULTS


Campylobacter jejuni Outbreak

7-day moving averages




RESULTS Gl Outbreak 2

Top 5 ICD-9 codes used during a Salmonella outbreak




Top 5 prescriptions during the Salmonella outbreak



RESULTS

Salmonella Outbreak

7-day moving averages




Analgesics/antipyretics: Acetaminophen, Tylenol, etc. **Antidiarrheals**: Loperamide, Imodium, Pepto-bismol, etc.


Antihistamines: Includes Phenergan, Promethazine (antiemetics)

Quinolones: Ciprofloxacin, Levaquin, etc.

RESULTS GI OUTBREAK 3

- At a U.S. Air Force training base in Texas³
 - Norovirus outbreak
 - In all 464 trainees affected
 - Lasted 7 days
 - 7/7 stool samples tested positive for Norovirus






RESULTS

Norovirus Outbreak

7-day moving averages




RESPOutbreaks


RESULTS RESP OUTBREAK 1

- At the Marine Corps Recruit Depot in San Diego³
 - Largest Group A Streptococcus outbreak since 1968
 - 160 recruits admitted for pneumonia (radiographically confirmed)
 - November 1 December 20
 - Sputum, blood and throat cultures
 - Group A Streptococcus isolated

Top 5 ICD-9 codes used during a Streptococcus A outbreak




Top 5 prescriptions written during a Streptococcus A outbreak



RESULTS

Streptococcus A Outbreak

7-day moving averages





Expectorants: Robitussin, Guaifenesin, etc.

Sympathomimetic agents: Sudafed, etc

Zinc replacement: Cepacol Cold Care, etc.


Analgesics/antipyretics: Acetaminophen, Tylenol, etc.

RESULTS RESP OUTBREAK 2


- Influenza in an Advanced Individual Training population in VA in November, 2003
 - Approx. 187 cases total
 - Oct. 31^{st} Nov. 7^{th}
 - Rapid flu tests positive; 5 CDC viral cultures Flu A positive, subtyped Korea-like or Fujian-like viruses

RESULTS RESP Outbreak

Top 5 ICD-9 codes used during an influenza outbreak




Top 5 prescriptions written during an influenza outbreak



RESULTS

Influenza Outbreak

7-day moving averages





RESULTS

GI group

- 1. Antiemetics
- 2. Antidiarrheals
- 3. Antihistamines
- 4. Analgesics/antipyretics -
- 5. Antispasmodics
- 6. Antacids
- 7. Quinolones
- 8. Intestinal adsorbents and protectives
- 9. Penicillins
- 10. Absorbable sulfonamides

RESP group

- 1. Expectorants
- Cough and/or cold preparations
- 3. Analgesics/antipyretics
- 4. Antitussives, non-narcotic
- 5. Sympathomimetic agents
- 6. NSAIDS
- 7. Macrolides
- 8. Quinolones
- 9. Zinc replacement
- 10. Penicillins

STUDY OBJECTIVES


- Determine medications commonly prescribed for GI and Respiratory (RESP) syndromes
- 2) Examine trends in daily counts of medications for GI, RESP, Asthma visits during outbreaks
- 3) Conduct retrospective surveillance on GI and RESP syndrome drug groups and GI and RESP outpatient visits at installations where outbreaks occurred

METHODS

- Retrospective surveillance conducted on pharmacy and outpatient visit data for approx. 1 year prior to and including the outbreak timeframe
 - Regression/Exponential Weighted Moving Average (EWMA) detection algorithms run on daily total GI/RESP visits, daily total PDTS syndrome groups and each GC3 group → output of p-values
 - P-values from individual larger count GC3 groups (>1 prescription in 3 days) were combined using Fisher and Edgington combination p-value methods
 - More information on combination p-value methods → Dr. Howard Burkom's talk at 2:00 p.m.
 - Detections with approximately 1 per 6 weeks false alert rate were identified graphically


Salmonella Outbreak

(12/9/02 to 12/18/02)




C. jejuni Outbreak

(11/16/02 - 11/18/02)




Norovirus Outbreak³

(02/17/04 to 02/24/04)




Norovirus Outbreak

(11/25/02 to 12/15/02)




Influenza Outbreak

(11/1/03 to 11/18/03)



Group A Streptococcus Outbreak

12/08/02 to 12/12/02



CONCLUSIONS

- Use of the GC3 classification system in military pharmacy data can increase sensitivity without significantly affecting the false alert rate.
 - A limitation: Certain medications were found in more than one GC3 group. If these don't contribute to an outbreak, the signal is weakened
- Outbreaks were characterized by increases in certain GC3 groups and combinations of GC3 groups but these varied by outbreak type.
- Syndromic drug groups were formed using medications more commonly used during outbreaks, increasing the overall sensitivity of the system.

CONCLUSIONS

- The ESSENCE outpatient visit detector, the gold standard, alerted for all 6 outbreaks.
- The ESSENCE pharmacy detector performed well and alerted for 5/6 outbreaks.
- The two p-value combination methods have distinct advantages. The Fisher method alerted for 5/6 outbreaks and was more sensitive to individual GC3 increases.
 Whereas, the Edgington method alerted for 3/6 outbreaks but was more responsive to consensus and in 2 outbreaks alerted earlier.
- Combination methods are more sensitive and specific if low count gc3 streams are accumulated or dropped.

FUTURE RESEARCH

- Investigate performance of algorithms for other outbreaks
- Investigate performance of detectors for a GC3 group switch, i.e., ignore product groups that don't meet a certain minimum count
- Continue to investigate product groups that give more signal and less noise
- Prospective surveillance using ESSENCE pharmacy and combination method detectors

REFERENCES

- 1. http://www.pec.ha.osd.mil/ICD/ICD_MTF_040502.doc
- Centers for Disease Control and Prevention.
 Outbreak of group A streptococcal pneumonia among Marine Corps recruits California, November 1 December 20, 2002. MMWR. 2003;52(6):106-109.
- 3. Yamane, GK, Shibukawa-Kent, RL. 2004. *A large norovirus GI outbreak among trainees at a USAF training base.* Seventh Annual Force Health Protection Conference, Aug. 8-12, 2004.

ACKNOWLEDGEMENTS

CPT Jeffrey K. Brown, USAMRID, Fort Detrick, MD

LCDR Scott Thornton, NEPMU-6, Pearl Harbor, HI

MAJ Samuel Jang, CHPPM, Aberdeen Proving Grounds, MD

Asha Riegodedios, MPH, NEHC, Bethesda, MD

THANK YOU FOR YOUR ATTENTION